

On the Production of Jets and Hadrons

S.Mukherjee(nee Banerjee) [†]

School of Mathematical Sciences,
Dublin Institute of Technology,
Dublin,Ireland.

Abstract

The ratio of the semivertical cone angles of the quark and gluon jets calculated in the statistical model is found to be in exact agreement with that derived from QCD. In hadron production also, the model's prediction of the power law type of behaviour for the inclusive distribution is strikingly close to the corresponding experimental findings; these spectacular results are significant for the model to be realistic.

PACS: 13.87.Ce, 13.87.Fh, 13.85.Ni, 13.66.Bc

[†] E-mail: smukherjee@dit.ie

1.Introduction

It is wellknown that colour forces organise quarks and gluons into hadrons through the fragmentic process. Becasue of the non-perturbative processes dominating it, exact theoretical prediction is still lacking and only predictions of modeltype calculations are available. A characteristic property of many of the most interesting events in hadron collider data is the appearance of jets which are dominated by shortdistance effects so that the perturbative Quantum Chromodynamics(pQCD) may be safely applied to study jet processes. As the hard quark scattering is manifest through the jet production; it becomes essential to study its inclusive production and to analyse the scattering at the largest possible transverse momenta(p_t)scale possible. In high energy pp interactions, the measurements of π^0 meson spectra in a high p_t ($p_t > 2GeV/c$) domain show a strong deviation from the exponential shape and Gazdzicki et. al[1] have found that there is the power law in hadron production and the mean multiplicity and transverse mass spectra obey such a law. Moreover, the estimate of the parameters in the power law are found to be the same when these are adjusted against corresponding experimental data for the production of different mesons at fixed collision energy. The results of our investigations on the jet observables, cross sections for particle production etc. have been compared with the relevant experimental findings and the predictions of pQCD, which help us to analyse the current status of the Statistical Model.

The Statistical Model of the hadron as a quarkonium system was proposed more than a couple of decades ago[2] and has

been widely used to study the properties of hadrons. Since, the model has also undergone modifications [3] along with its various ramifications and applications [3,4]. In the original version of the model [2] to be referred hereafter as the model (A), no allowance has been made for the exchange effect and we come across the fractal dimension $D = \frac{9}{2}$ of the hadron. However, for comparatively moderate values of the momenta of the constituents of a hadron so that the exchange energy effect may play an appreciable role [2], we arrive at $D=8.5$ in the modified version of the model (B). In the current investigation, the ratio $\frac{9}{4}$ expressed in terms of the angular radii of the quark and gluon jets, has been derived only from the geometric properties of the hadron in the framework of the Statistical Model (A). Moreover, the analysis of the cross sections for production of jets and hadrons reveals the power law falling of as p_t^{-D} where p_t is the transverse momentum of the secondary and this is found to be in accord with the experimental findings for moderate as well as for asymptotically large values of p_t .

It may be relevant to recall that the boundary of a jet reveals a hierarchy of indentations with a complex local structure and it behaves like a fractal [5] and the concept of fractal trees was imbibed to model the jets that arise when particles collide head on at very high energy. Hence it would be interesting to study if such a behaviour also exists in hadronic jets as the differences between the quark and gluon jets are also known to originate from QCD and it is curious to know if such a pattern is also manifest through the geometrical properties. Since the quarks and

gluons are viewed as carriers of colour charges, the asymptotic prediction of the ratio as that of the average parton multiplies in jets can thus be related to the charged hadron multiplicities [6]. However, without using the assumption of the parton model, Sterman and Weinberg [7] have investigated the properties of hadronic jets in e^+e^- annihilation in QCD and have asserted that their properties follow as consequences of pQCD and thus Einhorn and Weeks [8] have used QCD to investigate the properties of jets produced by a gluon source and have derived the relation between the jet opening angles. Shiuya and Tye [9] have also calculated the jet angular radius of a gluon jet in the framework of the pQCD and have estimated the ratio of the quark jet angular radius (δ_q) to that of the gluon jet (δ_g). Defining the fraction f of all jet like events which have a fraction $(1 - \epsilon')$ of the total energy E inside some pair of opposite cones of half angle δ we have,

$$f(gluon) = 1 - \frac{\alpha(E)}{\pi} [4C_2(G) \ln 2\epsilon' + \frac{11}{3}C_2(G) - \frac{4}{3}N_f T(R)] \ln \delta \quad (1)$$

The same formula for the quark case [7] is

$$f(quark) = 1 - \frac{\alpha(E)}{\pi} [4 \ln 2\epsilon' + 3] C_2(R) \ln \delta \quad (2)$$

where the colour factor $C_2(R) = \frac{4}{3}$, $C_2(G) = 3$ and $T(R) = \frac{1}{2}$ for a quark triplet and all other symbols have their usual meanings. For the same values of f, E, ϵ' , the ratio of the angular radii of the quark and gluon jets is independent of the coupling constant $\alpha(E)$ and in the asymptotic limit of p^2 , the first order prediction

for the jet angular sizes yields the ratio[8,9] as

$$\frac{\ln\delta_q}{\ln\delta_g} \rightarrow \frac{C_2(G)}{C_2(R)} = \frac{9}{4} \quad (3)$$

where δ_q and δ_g are in radians. On the other hand, in our model for the quark jet, we have a cone of semivertical angle δ_q where the volume of the right circular cone dimensions is $\frac{\pi h^3}{3} \tan^2 \delta_q$ where h is the height of the cone. Approximating $\tan \delta_q$ by δ_q for δ_q very small, we get $\delta_q^2 = \frac{3V}{\pi h^3}$ and hence in the context of the model (A), we have in the D dimensional space of the hadron with β a constant factor, $\delta_q^2 = \frac{\beta V}{h^D}$ and we get

$$\frac{\ln\delta_q}{\ln\delta_g} = \frac{\ln\delta_q^2}{\ln\delta_g^2} = \frac{\ln(\beta V)/(h^D)}{2\ln\delta_g} = \frac{1}{2} \left[\frac{\ln(\beta V/h^D)}{\ln\delta_g} \right] \quad (4)$$

As in the boxcounting method, we consider the quark jet to be covered with N balls of radius r and $\frac{r}{h} = \epsilon$, $\frac{V}{h^D} \sim \epsilon^D$, the unit of measurement where ϵ is a dimensionless very small quantity ($\epsilon \ll 1$) which is the size of the covering balls normalised by the linear size of the structure. Hence we have $N(\epsilon) \sim \epsilon^{-D}$, the number of D dimensional balls of radius ϵh needed to cover the fractal dimension D defined as $\lim_{\epsilon \rightarrow 0} \left[\frac{\ln N(\epsilon)}{\ln \epsilon^{-1}} \right]$. From (4) we have therefore,

$$\frac{\ln\delta_q}{\ln\delta_g} = \frac{1}{2} \frac{\ln(\beta V/h^D)}{\ln(r/h)} \rightarrow \frac{1}{2} \lim_{\epsilon \rightarrow 0} \frac{\ln \epsilon^D}{\ln \epsilon} = \frac{1}{2} \lim_{\epsilon \rightarrow 0} \frac{\ln N(\epsilon)}{\ln \epsilon^{-1}} = \frac{D}{2} = \frac{9}{4} \quad (5)$$

Hence we have

$$\delta_g = (\delta_q)^{4/9} \quad (6)$$

This remarkable result was also obtained by Einhorn et al.[8] and Shizuya et al[9] from QCD. Thus the aforesaid relation(6)

which indicates that the gluon jet spreads more than the quark jet has been derived in the current investigation purely from (fractal)geometrical properties of the hadron suggested by the model.

As the power law behaviour can often be attributed to the consequences of the underlying selfsimilarity,we analyse its predictions of the decrease of the crosssections of pp, $p\bar{p}$ collisions and e^+e^- annihilation in p_t in the context of the model.The analysis of these experimental data for the production of particles is likely to provide the information required to make important and stringent tests of its internal consistency.Thus,recently we [10] have derived and analysed the power law scaling in the light of the model and have found that it works well with $\alpha = 8.5$ as an input corresponding to the model (B) for the transverse mass spectra (m_t) of pions ($m_t > 1\text{Gev}/c^2$) and for particles from eta to upsilon.Our result are found to be in excellent agreement with the corresponding experimental findings with $D = 8.5$ for $p_t > 2\text{Gev}/c$. On the other hand,Gazdzicki et al[1] have indicated that the properly normalised multiplicities and transverse mass (m_t) spectra of neutral mesons obey the scaling law $cm_t^{-\alpha}$ where $m_t = \sqrt{m^2 + p_t^2}$ and c and α are similar for all mesons produced at the same collision energy.Introducing the hadronic mass spectrum)such that $\rho(m_t)$ such that $\rho(m_t)dm_t$ is the number of different hadron states in the transverse mass interval (m_t, dm_t) , we have for

$$\rho(m_t) \sim p_t^{-\alpha} \quad (7)$$

for $p_t \gg m$.Our assertion is that the crosssection for the pro-

duction of hadrons is proportional to the hadronic mass spectrum with $\alpha = 8.5$ and 4.5 for moderate values and asymptotic large values of p_t respectively. We would like to assert, as in Hagedorn[11], that almost all strong interactions may be simulated through $\rho(m_t)$ as the highly excited hadronic matter or fireball contains undetermined number of all species of hadrons and as such the scaling property displayed in (57) is also expected for jets observed in hadron collisions, jet fragmentation etc. On the other hand, the inclusive production of pions with large p_t in high energy pp collisions has been extensively studied and the invariant cross sections are also observed to obey a scaling law of the form

$$E \frac{d\sigma}{d^3p} \propto y(p_t = p_t^{-\alpha} I(x_t, \theta)) \quad (8)$$

where E and p are the energy and momentum of the secondary, θ is the production angle, $p_t = p \sin \theta$ is the transverse momentum, the scaling variable $x_t = \frac{2p_t}{\sqrt{s}}$ and y is a scaling function of p_t . In the case of pure hadronic collisions, the inclusive cross sections of the production of secondaries having limited p_t with respect to the collision axis possess the Feynman scaling and leads to the asymptotic behaviour with high p_t in $y(p_t)$ as p_t^{-4} in contradiction with available experimental data. However, to match the quark counting rule, the cross section must have the asymptotic p_t^{-8} (pion) if the production of high p_t particles is conditioned by hard scattering of a quark from a composite system of two quarks. Field and Feynman [12] have found that all experiments on production at 90° scales as in (8) with α lying between 7.2 and 8.6 . Feynman et al. [13] have also reexamined (8) for the large p_t

production of particles and jets in QCD approach for $pp \rightarrow \pi + X$ at $\theta_{cm} = 90^\circ$ for some moderate values of p_t and x_t and have found that $\alpha = 8$ and the right hand side of (8) becomes a universal function of x_t and p_t only and the falling of the afore-said experimental results according to (8) is found to be close to our prediction(B) as $\alpha = 8.5$. However, for $p_t = 10 \text{ GeV}/c$ at $x_t = 0.2$, it is close to p_t^{-5} and this is again in reasonable accord with our prediction(A) i.e. $\alpha = 4.5$. Clark et al [14] have reported measurements of inclusive π^0 production from high energy pp collisions at very large p_t at Fermilab and at CERN ISR at the centre of mass energies 58 and 63 GeV and $\theta \simeq 90^\circ$ and have found that the data are in accord with (8) and that lies in the range of 7 to 9 for x_t between 0.2 and 0.45. Experimental findings on charged pions from Fermi lab by Antreasyan et al [15] with 200, 30 and 400 GeV pp collisions for p_t ranging from 0.77 to 6.91 GeV/c also suggest that lies α between 8.2 and 8.5. Again, Ellis et al [16] have pointed out that $\sqrt{s} = 30.8 \text{ GeV}$ at ISR, and $\alpha \simeq 9$ and the study by McCubbin [17] on hadron production for large p_t in deep inelastic scattering of hadron hadron process also reveals that when the invariant single particle cross section is plotted against large p_t for over the energy range from 5 to 63 GeV, α turns out to be equal to 8.3 and α becomes 8 when the energy ranges from 20 to 60 GeV. The steeper falling of the cross section for inclusive distribution up to 10 GeV/c at ISR energies behaves as p_t^{-8} which compares favourably with the model's prediction of $p_t^{-8.5}$. On the other hand, at much higher energies, the $p_t^{-4.5}$ type of asymptotic dependence of $y(p_t)$ is suggested by the model (A) and obviously it compares reasonably well with the correspond-

ing QCD of p_t^{-4} and the experimental findings. Further in the trigger bias effect, regarded as a very important consequence of scaling in the jet fragmentation, the associated distribution on the trigger side is also found to scale in the particle distribution $\frac{d\sigma}{dp_t}$ as proportional to p_t^{-9} compared to the model's prediction (B) of $p_t^{-8.5}$ when $\frac{d\sigma}{dp_t}$ is assumed to be proportional to $\rho(m_t)$.

In studying single particle cross sections of the process $pp \rightarrow \pi + X$ at $\theta_{cm} = 90^\circ$, Feynman et al [13] have observed its p_t^{-8} behaviour at $p_t \leq 6 \text{ GeV}/c$ at $x_t = 0.2$ and at $p_t \leq 10 \text{ GeV}/c$ at $x_t = 0.5$ along with its asymptotic approach like p_t^{-4} . We may assert, therefore, in view of the above data that $y(p_t)$ has the cross over point p_t^c of the p_t variable where $p_t^c = 6 \text{ GeV}/c$ or $10 \text{ GeV}/c$ for the two cases respectively. As different exponents (8.5 and 4.5) are needed on either side of p_t^c for both the cases as in a selfaffine curve, the invariance of the measure is recovered by the affine type of transformation as in selfaffine fractals. The ratio of the semivertical angles of the quark and gluon jets in the asymptotic limit coincides exactly with that predicted from QCD; moreover, the power law decrease of the inclusive cross sections as $p_t^{-8.5}$ in $pp, p\bar{p}$, collisions and in e^+e^- annihilation is consistent with the corresponding experimental findings. Further, for asymptotically large values of p_t , the aforesaid power law becomes $p_t^{-4.5}$ which is not only in accord with the QCD prediction but it also agrees with the corresponding experimental data which fall approximately as p_t^{-4} . The model credited with the successes of these striking predictions and agreements with the experimental findings vouches for its realistic character for the

description of nature.

Acknowledgements

The author is thankful to Prof.S.N.Banerjee,Department of Physics,Jadavpur University,Kolkata,India for encouragement.

References

- [1] M.Gazdzicki and M.I. Gorenstein,Phys.Lett.B 517,250(2001).
- [2] S.N.Banerjee et. al., Had. J.4,2003(1981);(E)5,2157(1982); 6, 440,956(1983);7,186,795(1984);J.Phys.G.8,L61(1982); Ann.Phys. (N.Y.)150,150(1983); Nuov.Cim.A102,1733(1989)
- [3]S.N. Banerjee et.al.,Had. J. 11,243(1988);12,179(1989);13,75(1990).
- [4] S.N.Banerjee et.al.Int. J.Mod. Phys.A10,201(2001); Int.J.Mod.Phys.A17,4939(2002).
- [5]B.Mandelbrot,The Fractal Geometry(W.H. Freeman and Company,New York,U.S.A;1983)98.
- [6]S.J.Brodsky and J.F. Gunion, Phys.Rev.Lett. 37,402(1976); Skluth,Rep.Prog.Phys. 65,1771(2006).
- [7] G.Sterman and S.Weinberg,Phys. Rev.Lett.39,1436(1977).
- [8]M.B.Einhorn and B.G.Weeks,Nucl.Phys.B146,455(1978).
- [9]K.Shizuya and S.H.H. Tye,Phys.Rev.lett 41,787(1978).
- [10] S.N.Banerjee and S.Banerjee,Phys.Lett.B644,45(2007).
- [11] R.Hagedorn,CERN 71-72,38(1971).
- [12] R.D.Field and R.P. Feynman,Phys.Rev.D15,2590(1977).
- [13]R.P.Feynman et al.,Phys. Rev.D18,3320(1978).
- [14] A.G. Clark et al,Phys.Lett.74B,267(1978)
- [15]D.Antreasyan et al.,Phys.Rev.Lett.38,112(1977).
- [16]S.D.Ellis et al.,Nucl.Phys.B108,93(1976).
- [17]N.A.McCubbin,Phys.Scr.25,143(1982).
- [18]G.Giacomelli and M.Jacob,Phys.Rep.55,82(1979).
- [19] M.D.Shapiro and J.L. Siegrist,Ann.Rev.Nucl.Part.Sci,41,97(1991).